

AD-A114 467

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/O 12/1

A NEW CLASS OF HERMITE-BIRKHOFF QUADRATURE FORMULAS OF GAUSSIAN--ETC(U)

DEC 81 N DYN

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2314

NL

1 of 1

of 12/1

1

END

DATE

FORMED

6 82

DTIC

2

DA114487

MRC Technical Summary Report #2314

A NEW CLASS OF HERMITE-BIRKHOFF
QUADRATURE FORMULAS OF GAUSSIAN TYPE

Nira Dyn

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

December 1981

(Received September 24, 1981)

DTIC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

Approved for public release
Distribution unlimited

DTIC
ELECTE
MAY 18 1982
S E D

89 07 18 057

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

A NEW CLASS OF HERMITE-BIRKHOFF QUADRATURE FORMULAS
OF GAUSSIAN TYPE

Nira Dyn*

Technical Summary Report #2314
December 1981

ABSTRACT

It is shown how to combine incidence matrices, which admit Hermite-Birkhoff quadrature formulas of Gaussian type for any positive measure, in such a way that the resulting matrix also admits Gaussian type quadratures for any positive measure. Moreover, the uniqueness property and the extremal property of the formulas corresponding to the submatrices are transferred to the formula admitted by the composed matrix.

AMS (MOS) Subject Classifications: 41A05, 41A55

Key Words: Gaussian quadrature formulas, Hermite-Birkhoff interpolation

Work Unit Number 3 (Numerical Analysis and Computer Science)

*School of Mathematical Sciences, Tel-Aviv University, Israel.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

SIGNIFICANCE AND EXPLANATION

Methods for approximating the integral of a function, given the values of the function and/or some of its derivatives at several points in the interval of integration, are investigated. The integral is approximated by a weighted sum of the given data--a quadrature formula.

It is shown that for a wide class of different data configurations, there exist appropriate points of evaluation and weights such that the resulting quadrature formula is exact for all polynomials of the maximal possible degree. The well-known Gaussian quadrature formulas represent the particular case in which only function values (and no derivatives) are employed.



| | |
|--------------------|-------------------------------------|
| Accession For | |
| NTIS GRA&I | <input checked="" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By _____ | |
| Distribution/ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A | |

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

A NEW CLASS OF HERMITE-BIRKHOFF QUADRATURE FORMULAS OF GAUSSIAN TYPE

Nira Dym*

1. Introduction

Recently results concerning the existence of quadrature formulas of Gaussian type related to Hermite-Birkhoff interpolation problems have been obtained by several authors [5], [1], [2], [3]. Given the incidence matrix $E = \{e_{ij}\}_{i=0, j=0}^{m+1, n-1}$ with entries consisting of zeroes and ones, a Hermite-Birkhoff quadrature formula of Gaussian type (HB-QGF) is defined as a formula of the form

$$(1.1) \quad \int_a^b p d\sigma = \sum_{e_{ij}=1} a_{ij} p^{(j)}(x_i), \quad p \in \Pi_{n-1},$$

such that the number of parameters in (1.1) equals the dimension of Π_{n-1} (the space of polynomials of degree $\leq n-1$):

$$(1.2) \quad n = \sum_{i=0}^{m+1} \sum_{j=0}^{n-1} e_{ij} + m.$$

In (1.1) $d\sigma$ is a non-negative measure supported on more than m points in (a, b) , $(x_0, \dots, x_{m+1}) \in S^m$ where

$$S^m = \{Y = (y_0, \dots, y_{m+1}) \mid a = y_0 < y_1 < \dots < y_m < y_{m+1} = b\}$$

and $\sum_{j=0}^{n-1} e_{ij} > 0, 1 \leq i \leq m, \sum_{j=0}^{n-1} e_{ij} > 0, i = 0 \text{ or } m+1.$

This notion of HB-QGF extends the classical notion of Gaussian quadrature formulas

(GQF):

$$\int_a^b p d\sigma = \sum_{i=1}^m a_i p(x_i), \quad p \in \Pi_{2m-1}$$

* School of Mathematical Sciences, Tel-Aviv University, Israel.

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

where $a < x_1 < \dots < x_m < b$ and $a_i > 0$, $i = 1, \dots, m$, and also the notion of the multiple nodes GQF [6], [7].

The problem of characterizing incidence matrices admitting HB-GQF is posed in [4], and the following necessary condition on such matrices is proved:

Result A. Let $E = \{e_{ij}\}_{i=0, j=0}^{m+1, n-1}$ admits a quadrature formula exact for Π_{n-1} , for some non-negative measure $d\sigma$ supported on more than m points of $[a, b]$, and let r be the minimal number of ones which must be added to E to obtain a matrix $\tilde{E} = \{\tilde{e}_{ij}\}$ without odd sequences in rows $1, \dots, m$. Then \tilde{E} is a Pólya matrix (the number of ones in any first l columns exceeds $l - 1$), and

$$(1.3) \quad n \leq \sum_{i=0}^{m+1} \sum_{j=0}^{n-1} e_{ij} + r.$$

In [2] the existence of two classes of HB-GQF satisfying (1.3) with equality, has been claimed. The first class is related to incidence matrices with rows $1, \dots, m$ consisting of odd Hermite sequences and even non-Hermite sequences. The existence of this class of HB-GQF is proved in [1].

The present paper is concerned with the proof of the existence of the second class of HB-GQF, corresponding to incidence matrices which can be decomposed vertically into several submatrices, each admitting an HB-GQF with equality in (1.3). It is also proved here that such a formula is unique and/or has an extremal property in case each of the HB-GQF corresponding to the submatrices is unique and/or has an extremal property of the following type:

Definition 1: The HB-GQF (1.1) related to $E = \{e_{ij}\}_{i=0, j=0}^{m+1, n-1}$ has an extremal property, if for any f satisfying $f^{(n)} > 0$ on $[a, b]$:

$$(1.4) \quad \min_{p \in P(E, f)} \left\{ (-1)^\mu \int_a^b (f - p) d\sigma \right\} = (-1)^\mu \int_a^b f d\sigma - \sum_{e_{ij}=1} a_{ij} f^{(j)}(x_i)$$

where $\mu = \sum_{j=0}^{n-1} e_{m+1, j}$ and

$P(E, f) = \{p | p \in \Pi_{n-1, p}^{(j)}(v_i) = f^{(j)}(v_i), \tilde{e}_{ij} = 1, \forall i \in S^m\},$
 with $\tilde{E} = \{\tilde{e}_{ij}\}$ the matrix of Result A related to E .

The existence of another interesting class of HB-GQF has been established in [3].
 These formulas are related to incidence matrices of pyramidal type:

(a) Each interior row, i , contains one sequence only which is of odd length ℓ_i starting at column k_i .

(b) There exist $1 < I < J < m$ such that
 $k_i + \ell_i > k_{i-1} > k_i, i = 2, \dots, I, k_i = 0, i = I, \dots, J, k_i + \ell_i > k_{i+1} > k_i, i = J, \dots, m-1$

(c) The two rows $i = 0, i = m+1$ have non-zero entries in arbitrary positions $k > k_1$ and $k > k_m$ respectively.

The proof of the existence in [3] is based on the extremal property of Definition 1, characterizing several other classes of HB-GQF, in particular the classical GQF and the multiple nodes GQF [6], [7]. This class of matrices, as well as the class in [1], can be used in composing new HB-GQF of the type discussed in Section 2.

2. Main Results

Given an incidence matrix $E = \{e_{ij}\}_{i=0, j=0}^{m+1, n-1}$ with n ones, and a set of points $X \in S^m$, we denote by $p(E, X, f)$ the interpolating polynomial from Π_{n-1} to f at the data (E, X) :

$$(2.1) \quad p(E, X, f)^{(j)}(x_i) = f^{(j)}(x_i), \quad e_{ij} = 1.$$

Lemma 1: Let E be a Pólya matrix with even sequences only in rows $1, \dots, m$, and let $\mu = \sum_{j=0}^{n-1} e_{m+1, j} > 0$. Then $[f - p(E, X, f)](-1)^\mu > 0$ on $[a, b]$ whenever $f^{(n)} > 0$ on $[a, b]$.

Proof: It is sufficient to consider functions satisfying $f^{(n)} > 0$ on $[a, b]$. The case $f^{(n)} > 0$ on $[a, b]$ is obtained by taking the limit of such functions.

Since E has only even sequences in rows $1, \dots, m$, all the Rolle's zeroes of the derivatives of $g = f - p(E, X, f)$ do not coincide with those prescribed by E . This together with the Pólya conditions

$$(2.2) \quad \sum_{j=0}^{k-1} \sum_{i=0}^{m+1} e_{ij} > k, \quad 1 \leq k \leq n-1, \quad \sum_{j=0}^{n-1} \sum_{i=0}^{m+1} e_{ij} = n$$

imply that the numbers of Rolle's zeroes of $g^{(k)}$ is at least $\left[\sum_{j=0}^{k-1} \left(\sum_{i=0}^{m+1} e_{ij} - 1 \right) \right]$. Therefore $g^{(n-1)}$ has at least one $\left(\sum_{j=0}^{n-1} \sum_{i=0}^{m+1} e_{ij} - (n-1) = 1 \right)$ zero, which is the maximal possible number since $g^{(n)} > 0$. Thus the zeroes of g and its derivatives are only the Rolle's extension of (E, X) , namely those prescribed by (E, X) and the corresponding Rolle's zeroes. In particular all the interior zeroes of g are of even multiplicities, and therefore g does not change sign in $[a, b]$.

To determine the sign of g , observe that if $e_{m+1, k} = 0$ then the right most zero of $g^{(k)}$ in $[a, b]$, δ_k , is equal or greater than the right most change of sign of $g^{(k+1)}$ in (a, b) , ϵ_{k+1} . Therefore $g^{(k)}(x)g^{(k+1)}(x) > 0$ in $(b - \delta_k, b)$, while if $g^{(k)}(b) = 0$ ($e_{m+1, k} = 1$) then $g^{(k)}(x)g^{(k+1)}(x) < 0$ in $(b - \epsilon_{k+1}, b)$. This together with $g^{(n)} > 0$ implies that $(-1)^\mu g(x) > 0$ for $x \in [a, b]$.

As a direct consequence of Lemma 1, we obtain:

Corollary 1: Under the conditions of Lemma 1, the Peano kernel $K(t, x) \equiv K(t, x|E, X)$ for the error in the interpolation prescribed by (E, X) :

$$(2.3) \quad f(x) - p(E, X, f)(x) = \int_a^b f^{(n)}(t) K(t, x|E, X) dt$$

satisfies

$$(2.4) \quad (-1)^n K(t, x) > 0, \quad (t, x) \in [a, b]^2.$$

With these preliminary results the existence of HB-GQF corresponding to vertically decomposable matrices can be obtained.

Theorem 1: Let $E = \{e_{ij}\}_{i=0, j=0}^{m+1, n-1}$ be an incidence matrix which can be vertically decomposed in l submatrices, each consisting of m_i rows and n_i columns respectively:

$$(2.5) \quad E = E(n_1) \oplus E(n_2) \oplus \dots \oplus E(n_l), \quad \sum_{i=1}^l n_i = n,$$

such that all the non-zero entries of row i , $1 \leq i \leq m$, belong to one of the submatrices only. If each submatrix $E(n_i)$ admits a HB-GQF exact for Π_{n_i-1} , for any positive measure, and if each $E(n_i)$ satisfies Result A with equality in (1.3), then E admits a HB-GQF exact for Π_{n-1} , for any positive measure, with nodes $X = (x_0, \dots, x_{m+1}) \in \Omega^m$, where

$$(2.6) \quad \Omega^m = \{X = (X^{(1)} \cup X^{(2)} \cup \dots \cup X^{(l)}) | X^{(j)} \in S^{m_j}, 1 \leq j \leq l\}.$$

Proof: It is sufficient to consider the case $l = 2$. Let $\tilde{E} = \tilde{E}(n_1) \oplus \tilde{E}(n_2)$ be related to $E = E(n_1) \oplus E(n_2)$ by Result A, let $Y \in S^{m_1}$ be the nodes of the HB-GQF for $E(n_1)$ and the measure $d\sigma$, and let $Z \in S^{m_2}$ be the nodes of the HB-GQF for $E(n_2)$ and the measure $\omega(t)dt$ with

$$(2.7) \quad \omega(t) = \int_a^b d\sigma(x) K(t, x|\tilde{E}(n_1), Y).$$

That $\omega(t)$ is of constant sign follows from Corollary 1.

Consider now the set of fundamental polynomials for interpolation at (\tilde{E}, X) with $X = Y^{(1)} \cup Z^{(1)}$:

$$(2.8) \quad q_{ij}^{(k)}(x_v) = \delta_{iv} \delta_{jk}, \quad \tilde{e}_{kv} = 1.$$

Then

$$(2.9) \quad p(\tilde{E}, X, f)(x) = \sum_{\tilde{e}_{ij}=1} f^{(j)}(x_i) q_{ij}(x)$$

and for any $p \in \Pi_{n-1}$

$$(2.10) \quad p(x) = \sum_{\tilde{e}_{ij}=1} p^{(j)}(x_i) q_{ij}(x),$$

$$(2.11) \quad \int_a^b p(x) d\sigma(x) = \sum_{\tilde{e}_{ij}=1} p^{(j)}(x_i) \int_a^b q_{ij}(x) d\sigma(x).$$

Now $\{q_{ij} | \tilde{e}_{ij} = 1, j < n_1\}$ is a basis for Π_{n_1-1} , since $q_{ij}^{(n_1)} \in \Pi_{n_2-1}$ vanishes on $(\tilde{E}(n_2), Z)$, and therefore is identically zero. Using the HB-GQF with nodes $Y \in S^{m_1}$ and coefficients $\{a_{ij} | e_{ij} = 1, j < n_1\}$ admitted by $E(n_1)$, which is exact for Π_{n_1-1} , we obtain:

$$(2.12) \quad \int_a^b q_{ij}(x) d\sigma(x) = \sum_{\substack{e_{kv}=1 \\ v < n_1}} a_{kv} q_{ij}^{(v)}(y_k) = 0 \text{ if } e_{ij} = 0 \text{ and } \tilde{e}_{ij} = 1, 0 < j < n_1.$$

Consider next the set $\{q_{ij} | \tilde{e}_{ij} = 1, n_1 < j < n\}$, consisting of n_2 polynomials in Π_{n-1} vanishing on $(\tilde{E}(n_1), Y)$. These polynomials can be represented as

$$(2.13) \quad q_{ij}(x) = \int_a^b q_{ij}^{(n_1)}(t) K(t, x | \tilde{E}(n_1), Y) dt.$$

Hence

$$(2.14) \quad \int_a^b q_{ij}(x) d\sigma(x) = \int_a^b dt q_{ij}^{(n_1)}(t) \int_a^b K(t, x | \tilde{E}(n_1), Y) d\sigma(x) = \int_a^b q_{ij}^{(n_1)}(t) \omega(t) dt.$$

But $q_{ij}^{(n_1)} \in \Pi_{n_2-1}$ for $e_{ij} = 1$, $n_1 \leq j < n$, and therefore the application of the HB-GQF admitted by $E(n_2)$ for the measure $\omega(t)dt$, with nodes $z \in S^{m_2}$ and coefficients

$\{b_{ij} | e_{i,j+n_1} = 1, 0 \leq j < n_2\}$, yields

$$(2.15) \quad \int_a^b q_{ij}^{(n_1)} \omega(t)dt = \sum_{\substack{e_{kv}=1 \\ n_1 \leq v < n}} b_{k,v-n_1} q_{ij}^{(v)}(z_k) = 0 \text{ if } e_{ij} = 0 \text{ and } \tilde{e}_{ij} = 1, n_1 \leq j < n.$$

Combining (2.11) with (2.12) and (2.15) we conclude that

$$(2.16) \quad \int_a^b p d\sigma = \sum_{e_{ij}=1} \left[\int_a^b q_{ij}(x) d\sigma(x) \right] p^{(j)}(x_i), \quad p \in \Pi_{n-1}.$$

Using the same arguments it is easy to show that

Theorem 2: $E = E(n_1) \oplus \dots \oplus E(n_\ell)$ admits a HB-GQF at $X = X^{(1)} \cup \dots \cup X^{(\ell)}$ for the positive measure $d\sigma$, if and only if $E(n_i)$ admits a HB-GQF at $X^{(i)}$ corresponding to the positive measure $\omega_i(t)dt$ with

$$(2.17) \quad \omega_i(t) = \left| \int_a^b K(t, x | E(n_1) \oplus \dots \oplus E(n_{i-1}), X^{(1)} \cup \dots \cup X^{(i-1)}) d\sigma(x) \right|.$$

A direct conclusion of Theorem 2 is

Corollary 2: The HB-GQF admitted by $E = E(n_1) \oplus \dots \oplus E(n_\ell)$ is unique for any positive measure, if and only if the HB-GQF admitted by $E(n_i)$, $1 \leq i \leq \ell$, is unique for any positive measure.

Another property of the HB-GQF admitted by $E(n_i)$, $i = 1, \dots, \ell$, which is transferred to the HB-GQF admitted by $E(n_1) \oplus \dots \oplus E(n_\ell)$, is the extremal property. Note that (1.4) can be formulated also as

$$\min_{p \in P(E, f)} [(-1)^\mu \int_a^b (f - p) d\sigma] = (-1)^\mu \int_a^b [f - p(\tilde{E}, X, f)] d\sigma.$$

Theorem 3: Let E satisfy the conditions of Theorem 1. If for any positive measure each $E(n_k)$, $1 \leq k \leq l$, admits a HB-GQF of the form (1.1) with the extremal property of Definition 1, then for any positive measure E admits a HB-GQF with the extremal property of Definition 1, but with Ω^m replacing S^m in the definition of $P(E, f)$.

Proof: It is sufficient to consider the case $l = 2$. Let $f^{(n)} > 0$, let $\xi = \xi^{(1)} \cup \xi^{(2)}$ be arbitrary in Ω^m , and define $g_1 = f - p(\tilde{E}, \xi, f)$, where \tilde{E} is related to E by Result A. Since $g_1^{(n_1)}$ is the error in the interpolation of $f^{(n_1)}$ at $(\tilde{E}(n_2), \xi^{(2)})$, it follows from Corollary 1 that

$$\begin{aligned} g_1^{(n_1)}(x) &= \int_a^b g_1^{(n_1+n_2)}(t) K(t, x | \tilde{E}(n_2), \xi^{(2)}) dt \\ &= (-1)^{\mu_2} \int_a^b f^{(n)}(t) |K(t, x | \tilde{E}(n_2), \xi^{(2)})| dt \end{aligned}$$

with $\mu_2 = \sum_{n_1 < j < n} e_{m+1, j}$, and therefore

$$(2.18) \quad (-1)^{\mu_2} g_1^{(n_1)}(x) > 0, \quad x \in [a, b].$$

Now $0 \in P(E(n_1), (-1)^{\mu_2} g_1)$, and hence by the extremal property of the HB-GQF admitted by $E(n_1)$ with nodes Y , we obtain

$$(2.19) \quad (-1)^{\mu_1} \int_a^b (-1)^{\mu_2} g_1 d\sigma > (-1)^{\mu_1} \int_a^b (-1)^{\mu_2} [g_1 - q_1] d\sigma$$

with $\mu_1 = \sum_{j=0}^{n_1-1} e_{m+1, j}$ and $q_1 = p(\tilde{E}(n_1), Y, g_1) \in \Pi_{n_1-1}$. In view of the definition of g_1 and q_1 , (2.19) becomes:

$$(2.20) \quad (-1)^{\mu} \int_a^b [f - p(\tilde{E}, \xi, f)] d\sigma > (-1)^{\mu} \int_a^b [f - p(\tilde{E}, Y \cup \xi^{(2)}, f)] d\sigma,$$

with $\mu = \mu_1 + \mu_2$. To complete the proof we use again Corollary 1 and conclude that

$$\begin{aligned}
& (-1)^\mu \int_a^b [f - p(\tilde{E}, Y \cup \xi^{(2)}, f)] d\sigma \\
(2.21) \quad & = (-1)^\mu \int_a^b \int_a^b [f - p(\tilde{E}, Y \cup \xi^{(2)}, f)]^{(n_1)}(t) K(t, x | \tilde{E}(n_1), Y) dt d\sigma(x) \\
& = (-1)^\mu \int_a^b \omega(t) [f - p(\tilde{E}, Y \cup \xi^{(2)}, f)]^{(n_1)}(t) dt
\end{aligned}$$

where $\omega(t)$ is defined by (2.7), and satisfy $(-1)^{\mu_1} \omega(t) > 0$ on $[a, b]$. Applying the extremal property of the HB-GQF, admitted by $E(n_2)$ for $(-1)^{\mu_1} \omega(t) dt$ at the nodes Z , we obtain for $g_2 = f^{(n_1)} - p^{(n_1)}(\tilde{E}, Y \cup \xi^{(2)}, f)$, satisfying $g_2^{(n_2)} = f^{(n)} > 0$, the inequality

$$\begin{aligned}
& (-1)^{\mu_2} \int_a^b [(-1)^{\mu_1} \omega(t)] [f^{(n_1)} - p^{(n_1)}(\tilde{E}, Y \cup \xi^{(2)}, f)] dt > \\
(2.22) \quad & > (-1)^{\mu_2} \int_a^b [(-1)^{\mu_1} \omega(t)] [f^{(n_1)} - q_2](t) dt
\end{aligned}$$

where $q_2 = p(\tilde{E}(n_2), Z, f^{(n_1)}) \in \Pi_{n_2-1}$. Observing that $[f - p(\tilde{E}, Y \cup Z, f)]^{(n_1)} = f^{(n_1)} - q_2$, we finally derive from (2.20), (2.21) and (2.22) that

$$\begin{aligned}
& (-1)^\mu \int_a^b [f - p(\tilde{E}, \xi, f)] d\sigma > (-1)^{\mu_2} \int_a^b [f - p(\tilde{E}, Y \cup Z, f)]^{(n_1)} (-1)^{\mu_1} \omega dt \\
& = (-1)^\mu \int_a^b [f - p(\tilde{E}, Y \cup Z, f)] d\sigma,
\end{aligned}$$

indicating the optimal property of the HB-GQF admitted by E at $Y \cup Z$.

The following two matrices are examples of matrices which admit HB-GQF in view of Theorem 1 and the results in [1] and [3]:

$$E_1 = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$E_2 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

The matrix E_2 admits a HB-GOF with the extremal property of Theorem 3, in view of the results in [3], [6] concerning the two submatrices of E_2 .

REFERENCES

1. Dyn, N., "On the existence of Hermite-Birkhoff quadrature formulas of Gaussian type", to appear in J. Approx. Theory. (Also Mathematics Research Center Technical Summary Report #1978, July 1979).
2. Dyn, N., "Hermite-Birkhoff quadrature formulas of Gaussian type", in "Approximation Theory III" (E. W. Cheney, Ed.), pp. 371-376, Academic Press, 1980.
3. Jetter, K., "A new type of Gaussian formulas", to appear in SIAM J. Num. Anal.
4. Lorentz, G. G., Riemenschneider, S. D., "Birkhoff quadrature matrices", in "Linear Spaces and Approximation Theory" (P. L. Butzer, B. Sz. Nagy, Eds.), pp. 359-373, ISNM, vol. 40, Birkhäuser-Verlag, Basel, 1978.
5. Micchelli, C. A., Rivlin, T. J., "Quadrature formulae and Hermite-Birkhoff interpolation", Adv. in Math. 11 (1973), 93-112.
6. Popoviciu, T., "Asupra anei generalizari a formulei de integrare numerica a lui Gauss (Roumanian)", Acad. Republ. Popul. Romine Studii Cerc. Mat. 6 (1955), 29-57.
7. Turán, P., "On the theory of mechanical quadrature", Acta Sci. Math. Szeged 12 (1950), par. A, 30-37.

ND/scr

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|--------------------------------------|---|
| 1. REPORT NUMBER 2314 | 2. GOVT ACCESSION NO. AD-A114 487 | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) A NEW CLASS OF HERMITE-BIRKHOFF QUADRATURE FORMULAS OF GAUSSIAN TYPE | | 5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period |
| | | 6. PERFORMING ORG. REPORT NUMBER |
| 7. AUTHOR(s) Nira Dyn | | 8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041 |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 3 - Numerical Analysis & Computer Science |
| 11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709 | | 12. REPORT DATE December 1981 |
| | | 13. NUMBER OF PAGES 11 |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 15. SECURITY CLASS. (of this report) UNCLASSIFIED |
| | | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Gaussian quadrature formulas Hermite-Birkhoff interpolation | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown how to combine incidence matrices, which admit Hermite-Birkhoff quadrature formulas of Gaussian type for any positive measure, in such a way that the resulting matrix also admits Gaussian type quadratures for any positive measure. Moreover, the uniqueness property and the extremal property of the formulas corresponding to the submatrices are transferred to the formula admitted by the composed matrix. | | |

